



# The Centennial Rotation

Cotton/Corn/Soybean Systems
Effects on Nutrient Uptake and
Removal in the Midsouth
(2004-2011)





### Introduction

Crop rotation has been used in farming systems for hundreds of years with modern rotations (green manures) begun as early as 1730 in England. The benefits of crop rotation in the south can be divided into three major areas that include: a) maintenance of crop yields; b) control of diseases, insects, and weeds; and c) prevention of soil erosion. Before the extensive use of chemical fertilizers, maintenance and/or improvement of yields were best achieved by improving the base fertility of the soil in which the crop was grown. This usually required growing a legume crop to promote nitrogen fixation or applying manure to provide additional organic nutrients. Corn/cotton rotations were used through the first three to four decades of the 20th century as animal power on the farm was extremely important and corn was needed as feedstock for the animals. Mechanization and inorganic fertilizer materials reduced the need for some crops, rotations decreased, and mono-crop agriculture gained in popularity. With today's farm policies and programs, and the freedom to choose different crop mixes, rotations are coming back into prominence. Field research across the cotton producing states supported crop rotation. However, growers were reluctant to rotate cotton because of government payments and crop rotations complicated production practices and presented extra challenges for producers.

The Mississippi Legislature authorized the establishment of an experiment station in the Yazoo and Mississippi Delta. This marked the beginning of research in the region and the Delta Branch Experiment Station which has now been in existent for more than 100 years. The station continues to meet the original objective of the experiment station and land-grant institution — that is to make agriculture a profitable enterprise. Early research in Mississippi included simple rotations and the use of manure on fields that had been used for cotton production. Mechanization shifted the agricultural industry from hand labor to machines and chemicals while today that shift continues with the introduction and acceptance of biotechnology. The shift from rotation to mono-cultural and gradually back to rotation brings us to the 21st century. Cotton, corn, soybean, grain sorghum, and rice production recorded record yields in recent years with the aid of new technology and advancements through research. Since 2001, cotton, corn, and soybean have had yields and record prices. Corn acreage has increased while cotton has decreased in response to profitability. Grain crops can be planted early. With irrigation, yield stability has led to shifts in the crop mix and some producers shifting from cotton altogether.

The purpose of this research project was to establish long-term rotations involving cotton, corn, and soybean with the crops to be grown with the most upto-date technology available. It was designed to examine the impact of rotations on the whole-farm enterprise while monitoring soil nutrients, nematodes, and other pests. Several cooperators were identified to assist in the overall management of the project in order to assure maximum utilization of the data collected.

### Research Objectives:

1. Determine the effects of long-term crop rotation with respect to yield and profitability while utilizing state-of-the-art technology.

2. Assess the impact of crop rotation on the whole-farm enterprise.

- 3. Monitor changes in soil nutrient status, nematode numbers and types, and weed species.
- 4. Demonstrate the long-term need for crop rotation for the next century

### Materials and Methods

The research study includes five crop rotation sequences along with

continuous cotton as the base systems. All crops in a rotation sequence are grown each season thus establishing 15 distinct 'treatments' that are replicated four times. The five crop rotation sequences include 1) corn-cotton, 2) corncotton-cotton, 3) corn-soybean, 4) soybean-corn-cotton, and 5) soybean-corncotton-cotton and are summarize in Table 1. Each plot contains eight 40-in rows 200 ft in length with a minimum of four rows harvested for yield determinations. Fertility requirements are determined from soil tests each year. All cultural practices are maintained as uniformly as possible taking into consideration the technology that is available. Plots are harvested with commercial equipment adapted for plot harvests. Each plot is sampled for nutrient status and soil acidity (liming). The nutrient management and pesticide regimen is selected based on the committee expertise and recommendations. Production inputs and returns are then analyzed to determine the overall effects of rotation on whole-farm economics. With the current systems, it will take 12 years for all rotation systems to cycle back to the same point and the sequences will repeat. The actual arrangement of the research field is shown in Figure 1.

## Davis R. Clark

Research Associate III

and

# M. Wayne Ebelhar

Research Professor and Agronomist

Mississippi State University

Delta Research and Extension Center

Stoneville, Mississippi






Table 1: Cropping sequence for long-term cotton-based rotation cropping system. All crops in each sequence to be grown each year. MAFES-DREC Stoneville, MS

MISSISSIPPI STATE

UNIVERSITY

	L DOTA	TION OT										
CENTENNIA	LROIA	TION 51	UDY									
	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Cyatam	2004	2003	3		5	6		8	9	10	11	
System	I		S	4	5	0	7	0	9	10	11	12
1	CT	CT	CT	CT	CT	CT	CT	CT	CT	CT	CT	СТ
·	O I	01	01	01	01	01	01	01	01	01	01	
2	СТ	CR	СТ	CR	СТ	CR	CT	CR	СТ	CR	СТ	CR
3	CR	CT	CR	CT	CR	CT	CR	CT	CR	CT	CR	СТ
4	CR	CT	CT	CR	CT	CT	CR	CT	CT	CR	CT	СТ
5	CT	CR	CT	CT	CR	CT	CT	CR	CT	CT	CR	СТ
6	CT	CT	CR	CT	CT	CR	CT	CT	CR	CT	CT	CR
7	CR	SB	CR	SB	CR	SB	CR	SB	CR	SB	CR	SB
8	SB	CR	SB	CR	SB	CR	SB	CR	SB	CR	SB	CR
9	SB	CR	СТ	SB	CR	СТ	SB	CR	CT	SB	CR	СТ
10	CT	SB	CR	CT	SB	CR	CT	SB	CR	CT	SB	CR
11	CR	CT	SB	CR	CT	SB	CR	CT	SB	CR	CT	SB
						-						
12	SB	CR	СТ	СТ	SB	CR	CT	CT	SB	CR	CT	СТ
13	СТ	SB	CR	СТ	CT	SB	CR	CT	CT	SB	CR	СТ
14	CT	CT	SB	CR	CT	CT	SB	CR	CT	CT	SB	CR
15	CR	CT	CT	SB	CR	CT	CT	SB	CR	CT	CT	SB
OT 0 11		00 0		00 0								
CT = Cotton		CR = Co	orn	SB = Sc	oybean							

Figure 2: Estimated nutrient uptake for specific crops based on selected yields.

### **Nutrient Uptake for Selected Crops**

Crop	Yield	N	Р	K	S
	bu or lb/A	.00000000000000000000000000000000000000	lb//	Α	
Corn	180	240	45	199	30
Soybean	60	314	26	170	20
Wheat	80	149	24	135	21
Cotton	1000	160	21	116	24
Rice	7000	112	26	139	12

To Convert P to P<sub>2</sub>O<sub>5</sub> multiply by 2.29 To Convert K to K<sub>2</sub>O multiply by 1.20

20

### Figure 1: Plot layout for Centennial Rotation, Delta Research and Extension Center, Stoneville, MS. Layout is specific for 2011 Cropping Season.




Figure 3: Estimated nutrient removal for specific crops based on selected yields.

### **Nutrient Removal for Selected Crops**

Crop	Yield	N	Р	K	S
	bu or lb/A		lb/ <i>F</i>	٠	
Corn	180	162	35	43	14
Soybean	60	240	21	71	6
Wheat	80	92	19	23	5
Cotton	1000	64	12	33	6
Rice	7000	70	19	23	6

To Convert P to P<sub>2</sub>O<sub>5</sub> multiply by 2.29 To Convert K to K<sub>2</sub>O multiply by 1.20



#### **Results and Discussion**

The first eight years of the Centennial Rotation program was completed in 2011 (100-yr rotation). Long-term cropping system rotations and long-term research are limited in their scope in many areas of the world or are no longer in existence. The Morrow plots at the University of Illinois and The Old Rotation at Auburn University are some of the oldest continuous plots in the US. In an effort to celebrate the centennial anniversary of the Delta Branch Experiment Station and a new era in agricultural technology, the Centennial Rotation was initiated in 2004 at the Delta Research and Extension Center at Stoneville, MS. The "treatments" as outlined in Table 1 show the first 12 years of the rotations and the crops being grown each year. The project was originally setup as a cottonbased system due the historic significance of cotton to this region of the United States. Only one system (treatments 7 and 8) does not contain cotton and is meant to document the long standing advantages of corn/soybean rotations. With recent shifts to grain production, this system has become quite important. The systems will not begin to repeat in the thirteenth season at which time some rotations will have completed six cycles, others four cycles, and the last system will have completed three cycles.

The summary of the first eight years of crop yields are shown in Table 2. Cotton yields in the continuous cotton area have the overall lowest yields for cotton compared to the other systems. The greatest cotton yields as expected, follow corn production. Insect pressure and adverse weather conditions in 2007 resulted in the lowest cotton yields to date. In that year cotton yields were at least 18.6% higher where some other crop had been rotated compared to the continuous cotton system. Over the years the range has been 13.1 to 41.8% higher yields (115.0 to 387.3 lb lint/acre) where cotton was in some rotation with corn compared to continuous cotton. Average cotton yields have varied across years ranging from 891.1 lb lint/acre in 2007 to a high of 1461.8 lb/acre. Corn yields in the same time frame have ranged from 192.3 to 212.0 bu/acre excluding 2011. the 2011 yields (91.2 bu/acre) were way below average due to a lack of irrigation in a timely fashion. Soybean yields have ranged from 50.3 to 78.5 bu/acre with the lowest yields in 2011 (Table 2). Weather problems such as hurricanes have caused some problems (lodging) but the yields have still been harvestable. Timely irrigation is a key to successful and consistent corn production as evident in 2011. Timing of the first irrigation is critical.

One of the areas of interest in the long-term rotation study deals with nutrient uptake and removal. Nitrogen (N), phosphorus (P), potassium (K), and sulfur (S) uptake and removal are being calculated for each of the systems. Figure 2 shows the estimated N, P, K, and S uptake for selected crops in the Mississippi Delta while Figure 3 gives an estimate of the N, P, K, and S removal by the crops based on the yields given. For cotton, corn, and soybean, the crops take up more nutrients than are actually removed from the field. Only the grain portion of corn and soybean are removed and the seed and lint portion of cotton along with some vegetative materials. Soybean removes the largest percentage of N and K while corn removes the largest percentage of P. These values have been used to calculate nutrient uptake and removal for the crop sequences that have been grown to date. The summary of nutrient uptake is shown in Table 3 and the summary of nutrient removal is shown in Table 4. As expected, the more cotton grown, the lower the N uptake and removal. The same is true for P and K also. The greatest N uptake and removal has occurred in the corn/soybean rotation system (Treatments 7 and 8). Much of the N that is removed in this system comes from symbiotic N fixation associated with soybean production and from high rates of fertilizer N addition for corn production. Producers should take extra steps to insure adequate fertility when shifting from cotton production to rotations with grain crops. Nutrient removal, especially N, can be 3 to 4 times higher than continuous cotton.

The economic impact of crop rotations is evident in most years just from the yield standpoint. However, as the costs of inputs continue to escalate, particularly with respect to technology fees, the more important rotation becomes. The increase in herbicide-resistant weed species across the country could lead to even more emphasis on crop rotation and herbicide rotation.





Table 3: Summary of total nutrient (N, P, K, S) uptake from the Centennial Rotation Study (2004 – 2011). Delta Research and Extension Center, Stoneville, Mississippi

NUTR	RIENT U	PTAKI							N	Р	K	S
	Crop Sequence							Uptake	Uptake	Uptake	Uptake	
Trt	2004	2005	2006	2007	2008	2009	2010	2011	(lb/acre)	(lb/acre)	(lb/acre)	(lb/acre)
1	СТ	СТ	СТ	СТ	СТ	CT	СТ	CT	1266.8	166.28	1266.83	190.00
2	CT	CR	CT	CR	CT	CR	CT	CR	1675.6	268.63	1675.63	229.69
3	CR	CT	CR	CT	CR	CT	CR	CT	1707.2	282.25	1707.17	230.18
4	CR	CT	CT	CR	CT	CT	CR	CT	1713.9	271.34	1713.87	236.48
5	CT	CR	CT	СТ	CR	CT	СТ	CR	1572.3	243.40	1572.28	219.39
6	CT	СТ	CR	СТ	CT	CR	CT	CT	1577.3	235.97	1577.29	223.73
7	CR	SB	CR	SB	CR	SB	CR	SB	2446.5	315.12	2446.54	221.73
8	SB	CR	SB	CR	SB	CR	SB	CR	2250.8	288.00	2250.83	202.86
9	SB	CR	CT	SB	CR	CT	SB	CR	2136.6	267.10	2136.59	209.30
10	CT	SB	CR	СТ	SB	CR	СТ	SB	2013.8	251.56	2013.81	211.90
11	CR	СТ	SB	CR	CT	SB	CR	СТ	2053.2	282.29	2053.23	228.54
12	SB	CR	CT	СТ	SB	CR	СТ	СТ	1798.4	235.79	1798.36	203.35
13	СТ	SB	CR	СТ	CT	SB	CR	СТ	1870.4	244.48	1870.40	212.12
14	CT	СТ	SB	CR	CT	CT	SB	CR	1853.9	236.28	1853.87	210.72
15	CR	СТ	СТ	SB	CR	CT	СТ	SB	1915.2	248.09	1915.15	214.52

Table 2: Summary of crop yields from the Centennial Rotation Study (2004 – 2011)

Delta Research and Extension Center, Stoneville, Mississippi

tation			Cr	op Yea	r				2004	2005	2006	2007	2008	2009	2010	2011
System	2004	2005	2006	2007	2008	2009	2010	2011	Crop	Crop	Crop	Crop	Crop	Crop	Crop	Crop
	1	2	3	4	5	6	7	8	Yield	Yield	Yield	Yield	Yield	Yield	Yield	Yield
1	СТ	СТ	СТ	СТ	СТ	СТ	СТ	СТ	1430.5	1101.8	978.9	718.5	927.6	877.6	1039.4	843.2
2	СТ	CR	СТ	CR	СТ	CR	СТ	CR	1470.9	204.6	1185.4	200.8	1218.9	182.4	1185.6	61.6
3	CR	CT	CR	CT	CR	CT	CR	CT	201.2	1334.3	185.1	942.2	194.9	961.3	194.7	965.4
Ü	OIX	01	OIX	O I	Oit		Oit		201.2	1004.0	100.1	042.2	104.0	001.0	104.7	000.1
4	CR	СТ	СТ	CR	СТ	СТ	CR	СТ	197.2	1298.4	988.0	219.4	1314.9	975.3	201.8	982.2
5	СТ	CR	CT	СТ	CR	СТ	СТ	CR	1509.4	213.3	1202.1	866.7	206.8	984.7	1148.2	73.8
6	CT	CT	CR	CT	CT	CR	CT	CT	1525.1	1148.8	191.1	909.3	982.5	194.8	1234.7	841.9
7	CR	SB	CR	SB	CR	SB	CR	SB	193.9	57.8	199.3	78.4	205.8	73.3	207.2	52.6
8	SB	CR	SB	CR	SB	CR	SB	CR	60.3	212.3	62.5	208.8	56.1	205.1	65.7	101.8
	0.5	0.0	0.7	0.5	0.5	<b>0</b> -	0.0	0.5	0.1.1	0400	10000		407.0	2215		440 =
9	SB	CR	CT	SB	CR	CT	SB	CR	61.4	212.6	1206.2	75.5	197.6	994.5	70.6	113.7
10	CT	SB	CR	CT	SB	CR	CT	SB	1447.5	61.5	194.6	1019.2	60.4	209.4	1199.0	47.9
11	CR	СТ	SB	CR	СТ	SB	CR	СТ	195.9	1268.2	64.4	207.6	1222.3	66.3	209.0	963.0
12	SB	CR	СТ	СТ	SB	CR	СТ	СТ	60.4	199.0	1152.6	852.2	57.5	195.9	1239.2	849.3
13	СТ	SB	CR	СТ	СТ	SB	CR	СТ	1402.7	52.3	191.2	929.5	978.7	69.8	208.0	1059.2
14	СТ	СТ	SB	CR	СТ	СТ	SB	CR	1446.6	1148.2	58.1	223.4	1240.5	929.3	66.8	105.0
15	CR	СТ	СТ	SB	CR	СТ	СТ	SB	200.5	1359.4	947.2	81.5	199.9	992.6	1026.1	50.4

Table 4: Summary of total nutrient (N, P, K, S) removal from the Centennial Rotation Study (2004 – 2011). Delta Research and Extension Center, Stoneville, Mississippi

MIITR	IENT R	<b>EMOV</b>	۸۱						N	Р	K	S
INOTIN				Crop Sequenc					Removal	Removal	Removal	Removal
Trt	2004	2005	2006	<del>ор Зеци</del> 2007	2008	2009	2010	2011	(lb/acre)	(lb/acre)	(lb/acre)	(lb/acre)
111	2004	2003	2000	2001	2000	2009	2010	2011	(ID/ACTE)	(ID/acre)	(ID/acre)	(ID/ACTE)
1	СТ	СТ	СТ	СТ	СТ	СТ	СТ	СТ	506.75	95.01	506.75	47.51
1	Ci	Ci	Ci	Ci	CI	CI	Ci	CI	300.73	95.01	500.75	47.51
2	СТ	CR	СТ	CR	СТ	CR	СТ	CR	908.28	187.01	908.28	80.87
3	CR	СТ	CR	СТ	CR	СТ	CR	СТ	967.33	201.31	967.33	85.57
4	CD	СТ	СТ	CD	СТ	СТ	CD	СТ	012.40	196.05	012.40	04 45
4	CR	CT	CT	CR	CT	CT	CR	CT	912.40	186.95	912.40	81.45
5	СТ	CR	CT	СТ	CR	СТ	СТ	CR	810.13	164.57	810.13	72.68
6	CT	CT	CR	CT	CT	CR	CT	СТ	772.42	154.75	772.42	69.87
7	CR	SB	CR	SB	CR	SB	CR	SB	1773.98	248.50	1773.98	88.92
8	SB	CR	SB	CR	SB	CR	SB	CR	1633.61	226.17	1633.61	81.08
9	SB	CR	CT	SB	CR	CT	SB	CR	1442.28	200.89	1442.28	74.71
10	CT	SB	CR	CT	SB	CR	CT	SB	1277.29	181.97	1277.29	70.41
11	CR	CT	SB	CR	СТ	SB	CR	CT	1108.06	197.76	1108.06	81.44
12	SB	CR	CT	СТ	SB	CR	CT	CT	1088.98	167.16	1088.98	67.07
13	CT	SB	CR	CT	CT	SB	CR	CT	1127.39	172.78	1127.39	69.49
14	СТ	СТ	SB	CR	СТ	СТ	SB	CR	1100.21	164.76	1100.21	66.63
15	CR	СТ	СТ	SB	CR	СТ	СТ	SB	1164.70	175.92	1164.70	70.29
									-	-		