

Zachary Eder, Gaylon Morgan, Daniel Fromme, Dale Mott, Fred Bourland, and John Kerns

Department of Soil and Crop Sciences  
Texas A&M University

## ABSTRACT

Defoliation of cotton, *Gossypium hirsutum* L., has been referred to as more art than a science by industry leaders. The remnants of leaf material in harvested cotton can significantly increase leaf grade values and result in dockage to the producer. Cotton classed through the USDA-AMS Classing office in Corpus Christi, Texas has reported increases in leaf grade values beginning in 2000. The impacts of the agronomic variables were studied during the 2010 and 2011 growing season and data collected were used to identify possible contributors to leaf grade, including leaf pubescence and harvest-aid treatments. Harvest-aid and harvest-aid by variety trials were initiated in 2010 and 2011. Variety by harvest-aid trials provided an approach to analyze the combined impact of the two factors. All samples were ginned on a miniature gin in Lubbock, HVI and fiber analysis was done. Wide ranges of percent defoliation and desiccation were obtained with the treatments, but had no significant impact on leaf grade during 2010 or 2011. In the variety by defoliation trial, the hairy variety of cotton had higher leaf grades than a smooth variety across multiple levels of defoliation in 2010 and 2011. Overall leaf grades were lower in 2011 due to more suitable weather conditions between harvest-aid application and harvesting.

## INTRODUCTION

Cotton leaf grade is a visual estimation of the amount of plant material in a lint sample on a scale of 1 to 7, with one being the ideal score (Larson and English, 2001). Plant material in harvest lint is waste, and can result in price dockage for the producer because additional processing is required to remove the plant material. Currently, several factors are believed to negatively influence the leaf grade values. First is the level of defoliation and desiccation prior to harvest. Second is the varietal characteristics, such as leaf hairiness, bract hairiness, and leaf and bract size.

The efficacy of chemical defoliation can be an unpredictable process but is vital for the harvest efficiency and to minimize dockage from plant materials (Valco and Snipes, 2001). Factors impacting defoliation vary from harvest-aid selection, plant condition, weather prior to and during application, spray coverage, canopy density, translocation, and varietal traits (Cathey, 1986, Oosterhuis et al. 1991). Additionally, hairier varieties are suspected of contribute to higher leaf grades through a "velcro effect". The hairiness of commercial cotton varieties are assigned by a subjective rating system (smooth to very-hairy), and inconsistencies exist between varietal ratings available to producers.

## HYPOTHESES

- Higher percentages of defoliation will cause a lower leaf grade in harvested cotton
- Reduction of desiccation will yield a lower leaf grade in harvested cotton
- Hairy leaf varieties of cotton will amplify leaf and bract trash resulting in higher leaf grades than smooth leaf varieties

## OBJECTIVES

- Identify the impact of traditional harvest-aid products on defoliation, desiccation and leaf grade
- Identify interaction for variety characteristics, defoliation and desiccation impacting cotton leaf grade

## MATERIALS AND METHODS

All trials were four rows wide and 40 feet long and were treated with a Lee Spider sprayer with 11 GPA carrier volume using XR flat fan tips. Percent defoliation, desiccation and green leaf were rated at 7 and 14 days after treatment. Untreated check was rated as zero and complete absence of leaves was 100%. Plots were mechanically harvested with a picker. Samples were ginned in a miniature gin, and leaf grade and fiber quality parameters were processed at the Fiber and Biopolymer Research Institute using HVI analysis.

In the defoliation trials, treatments were superimposed over a field of Phytogen 375WRF. Twenty defoliation treatments were applied to obtain a wide range of defoliation and desiccation levels in 2010 (Fig. 1 and 2). In 2011, sixteen defoliation treatments were selected to provide the range of defoliation and desiccation (Fig. 1 and 2).

In the variety by defoliation trials, Stoneville 5458B2RF, a hairy leaf variety, and DynaGro 2570B2RF, a smooth leaf variety, received five defoliation treatments and had four replications in a split block design. Treatments were applied to achieve variable defoliation levels. In 2010 the trial was conducted in Colorado county (Fig. 3), and in 2011 the trial was repeated in Matagorda county (Fig. 4).

ANOVA was performed and means separation using LSD with  $P=0.05$  (data not shown). Kruskal-Wallis was used to identify significance between leaf grades. Locations are shown separately due to significant location interaction.



Fig. 1. Cotton defoliant treatments vary in amount of defoliation and desiccation.




Fig. 2. Leaf grade of cotton in defoliation trials in four site-years based on defoliation ratings at 14 days after application of treatments.

\*Kruskal-Wallis test indicated the cotton leaf grade was not affected the defoliation treatment at any of the locations in 2010 or 2011 ( $P = 0.05$ ).




Fig. 3. Impact of leaf hairiness on the leaf grade of cotton treated with five harvest-aid treatments during the 2010 growing season.

\*Kruskal-Wallis test indicated the cotton leaf grade was affected by leaf hairiness ( $P = 0.05$ ).




Fig. 4. Impact of leaf hairiness on the leaf grade of cotton treated with five harvest-aid treatments during the 2011 growing season.

\*Kruskal-Wallis test indicated the cotton leaf grade was affected by leaf hairiness ( $P = 0.05$ ).

## RESULTS AND DISCUSSION

**Defoliation Trial:** A wide range of defoliation and desiccation levels were obtained with the selected defoliation treatments (Fig. 2). Despite the range of defoliation levels, no differences were observed in leaf grade values (Fig. 2). The 2010 season had leaf grades of 3 and 4, while in 2011 leaf grade values did not rise above 2. Low leaf grades in 2011 were the result of more suitable harvest conditions, compared to 2010.

**Variety Hairiness by Defoliation:** A good range of defoliation levels were obtained with the 5 defoliation treatments, and efficacy was comparable for the smooth leaf and hairy leaf varieties (Fig. 3). Leaf grade values were consistently lower across all defoliation levels for the smooth leaf variety (Fig. 3 and 4). Leaf grade ratings were less than 1.5 regardless of the defoliation level or variety hairiness (Fig. 4). Though 2011 conditions were suitable for low leaf grade, there was a variety affect on the scores (Fig. 4).

## CONCLUSIONS

- Cotton leaf grade was not influenced by the defoliation or desiccation levels
- Leaf hairiness influences leaf grade more than defoliation when environmental conditions are conducive for higher leaf grades
- Differences between years indicate specific environmental conditions, such as rainfall after harvest-aid application, needed for high leaf grade

## FURTHER RESEARCH

- Evaluate various physiological traits that may influences cotton leaf grades and other fiber quality characteristics
- Develop an industry-wide standard for leaf hairiness

## ACKNOWLEDGEMENTS

This research was supported by the Texas State Support Committee, Cotton Incorporated, Cotton Foundation, and Texas A&M University.

## REFERENCES CITED

- Cathey, G. W. 1986. Physiology of defoliation in cotton production. p. 143-154. In J. R. Mauney, and J. McD. Stewart (ed.) Cotton physiology. The Cotton Foundation Reference Book Series, Cotton Foundation, Memphis, TN.
- Larson, J. A. and B. C. English. 2001. Factors influencing net returns to cotton harvest aids. p. 181-206. In J.R. Supak and C.E. Snipes (eds) Cotton Harvest Management: Use and Influence of Harvest Aids. The Cotton Foundation Reference Book Series.
- Oosterhuis, D.M., R. E. Hampton, and S. D. Wullschlegar. 1991. Water deficit effects on the leaf cuticle and efficiency of defoliant. J. Prod. Agric. 4:260-265.
- Valco, T. D. and C. E. Snipes. 2001. Uniform harvest-aid performance and lint quality evaluation. p. 167-180. In J.R. Supak and C.E. Snipes (eds) Cotton Harvest Management: Use and Influence of Harvest Aids. The Cotton Foundation Reference Book Series.